



ОТКРЫТОЕ ПИСЬМО

Вч. Витебск.



Сашмуль Левицкому Гуревичу

в Д.Лейбс Гурвича зашкварь 16
премнг. Э. Ильинская 1 ДЕК.
1883



1. На этой сторонѣ кромѣ адреса не дозволяется ничего другаго писать.
2. Почтовое Управление за содержаніе письма не отвѣтствуетъ.

Drink

Water

Water can be given to the patient in the following ways:

- 1. Oral route: This is the easiest and most common way of giving water. It can be given in small sips or as a full glass. It is important to encourage the patient to drink as much water as possible.
- 2. Intravenous route: This is used when the patient is unable to drink orally due to nausea, vomiting, or unconsciousness. It involves inserting a needle into a vein and injecting the water directly into the bloodstream.
- 3. Nasogastric route: This is used when the patient has difficulty swallowing. A tube is inserted through the nose and into the stomach to deliver the water.
- 4. Rectal route: This is used when the patient is unconscious or has a closed airway. A rectal enema can be used to administer water into the rectum.

It is important to monitor the patient's fluid intake and output to ensure they are staying hydrated. If the patient is unable to drink, it may be necessary to provide fluids via a nasogastric or rectal route.

Water

Water can be given to the patient in the following ways:

- 1. Oral route: This is the easiest and most common way of giving water. It can be given in small sips or as a full glass. It is important to encourage the patient to drink as much water as possible.
- 2. Intravenous route: This is used when the patient is unable to drink orally due to nausea, vomiting, or unconsciousness. It involves inserting a needle into a vein and injecting the water directly into the bloodstream.
- 3. Nasogastric route: This is used when the patient has difficulty swallowing. A tube is inserted through the nose and into the stomach to deliver the water.
- 4. Rectal route: This is used when the patient is unconscious or has a closed airway. A rectal enema can be used to administer water into the rectum.

It is important to monitor the patient's fluid intake and output to ensure they are staying hydrated. If the patient is unable to drink, it may be necessary to provide fluids via a nasogastric or rectal route.